

Domain of a HASKELL-program

Least solution of

$$\text{Dom} = \text{Functions} \oplus \text{Tuples} \oplus \text{Constructions}_0 \oplus \text{Constructions}_1 \oplus \dots$$

where $\text{Functions} = \langle \text{Dom} \rightarrow \text{Dom} \rangle_{\perp}$

$$\text{Tuples} = \{\perp, ()\} \oplus (\text{Dom} \times \text{Dom})_{\perp} \oplus (\text{Dom} \times \text{Dom} \times \text{Dom})_{\perp} \oplus \dots$$

$$\text{Constructions}_n = (\text{Con}_n \times \text{Dom}^n)_{\perp}$$

Environment

$$\rho : \text{Var} \rightarrow \text{Dom} \text{ with } \rho = \{\underline{\text{var}}_1, \dots, \underline{\text{var}}_n / d_1, \dots, d_n\}$$

$$(\rho_1 + \rho_2)(\underline{\text{var}}) = \begin{cases} \rho_2(\underline{\text{var}}), & \text{if } \rho_2(\underline{\text{var}}) \text{ is defined} \\ \rho_1(\underline{\text{var}}), & \text{otherwise.} \end{cases}$$

Initial environment ω

for all pre-defined variables in HASKELL

Simple HASKELL-Programs

- no type synonyms, no type classes, no pre-defined lists
- only one declaration of the form var = exp

exp → var
| constr
| integer
| float
| char
| (exp₁, ..., exp_n), n ≥ 0
| (exp₁ exp₂)
| **if** exp₁ **then** exp₂ **else** exp₃
| **let** var = exp **in** exp
| \ var → exp

Semantics of HASKELL-Programs

- $\mathcal{V}al \llbracket \underline{\text{var}} \rrbracket \rho = \rho(\underline{\text{var}})$
- $\mathcal{V}al \llbracket \underline{\text{constr}}_0 \rrbracket \rho = \underline{\text{constr}}_0 \text{ in } \text{Constructions}_0 \text{ in Dom}$
- $\mathcal{V}al \llbracket \underline{\text{constr}}_n \rrbracket \rho = f \text{ in Functions in Dom, where } f d_1 d_2 \dots d_n = (\underline{\text{constr}}_n, d_1, \dots, d_n) \text{ in } \text{Constructions}_n \text{ in Dom}$
- $\mathcal{V}al \llbracket (\underline{\text{exp}}_1, \dots, \underline{\text{exp}}_n) \rrbracket \rho = (\mathcal{V}al \llbracket \underline{\text{exp}}_1 \rrbracket \rho, \dots, \mathcal{V}al \llbracket \underline{\text{exp}}_n \rrbracket \rho) \text{ in } \text{Tuples}_n \text{ in Dom,}$
where $n = 0$ or $n \geq 2$
- $\mathcal{V}al \llbracket (\underline{\text{exp}}) \rrbracket \rho = \mathcal{V}al \llbracket \underline{\text{exp}} \rrbracket \rho$
- $\mathcal{V}al \llbracket (\underline{\text{exp}}_1 \underline{\text{exp}}_2) \rrbracket \rho = f(\mathcal{V}al \llbracket \underline{\text{exp}}_2 \rrbracket \rho),$
where $\mathcal{V}al \llbracket \underline{\text{exp}}_1 \rrbracket \rho = f \text{ in Functions in Dom}$

$$\mathcal{V}al \llbracket \text{if } \underline{\text{exp}}_1 \text{ then } \underline{\text{exp}}_2 \text{ else } \underline{\text{exp}}_3 \rrbracket \rho = \begin{cases} \mathcal{V}al \llbracket \underline{\text{exp}}_2 \rrbracket \rho, & \text{if } \mathcal{V}al \llbracket \underline{\text{exp}}_1 \rrbracket \rho = \text{True} \\ & \text{in Constructions}_0 \text{ in Dom} \\ \mathcal{V}al \llbracket \underline{\text{exp}}_3 \rrbracket \rho, & \text{in } \mathcal{V}al \llbracket \underline{\text{exp}}_1 \rrbracket \rho = \text{False} \\ & \text{in Constructions}_0 \text{ in Dom} \\ \perp, & \text{otherwise} \end{cases}$$

$$\mathcal{V}al \llbracket \text{let } \underline{\text{var}} = \underline{\text{exp}}' \text{ in } \underline{\text{exp}} \rrbracket \rho = \mathcal{V}al \llbracket \underline{\text{exp}}' \rrbracket (\rho + \{\underline{\text{var}} / \text{lfp } f\}),$$

where $f : \text{Dom} \rightarrow \text{Dom}$ and $f(d) = \mathcal{V}al \llbracket \underline{\text{exp}} \rrbracket (\rho + \{\underline{\text{var}} / d\})$

$$\mathcal{V}al \llbracket \backslash \underline{\text{var}} \rightarrow \underline{\text{exp}} \rrbracket \rho = f \text{ in Functions in Dom} \\ \text{where } f(d) = \mathcal{V}al \llbracket \underline{\text{exp}} \rrbracket (\rho + \{\underline{\text{var}} / d\})$$